

Table 1: Superoxide production rates in *Synechococcus* high (PetC-Δ2G) and low (PetB-R214H) potential chain mutants

Strain	DCMU (@10 μM)	Superoxide Production	
		(pmol μg protein ⁻¹ h ⁻¹)	R ²
WT	–	60	0.9385
	+	40	0.9979
PetB R214H	–	100	0.9908
	+	90	0.8994
PetC Δ2G	–	50	0.9806
	+	30	0.9964

wild type showed a significant deuterium kinetic isotope effect on cytochrome *b* reduction/oxidation kinetics whereas D₂O had little effect on the mutant. Redox titrations of membranes revealed midpoint potentials (Em₇) of ~–30 mV and –120 mV for the two *b* hemes in both the mutant and wild type. These functional data define cytochrome *b*₆ arginine 214 as a key residue for Q_i-site function, cytochrome *bf* turnover, and photosynthetic growth. In the *bf* structures, R214 lies near the Q_i pocket and the newly discovered *c'* (or *x*) heme. Our data suggest involvement of PetB–Arg214 (*Chlamydomonas*, *Mastigocladus* R207) in plastoquinone binding, in midpoint potential determination of the *c'* (*x*) heme, and possibly in proton transfer reactions at the Q_i-site.

Production of reactive oxygen species (ROS) may result from blockage of electron flow through the low potential chain. To test this, we measured superoxide production in wild type, PetB–R214H, and PetC–Δ2G mutants of *Synechococcus* (Table 1). Mutations PetC–Δ2G in the high potential chain (Yan & Cramer 2003) and PetB–R214H in the low potential chain (current study) slow overall cytochrome *bf* turnover comparably. However, the R214H mutant generated superoxide at about twice the rate of wild type or PetC–Δ2G. Addition of DCMU did not significantly alter ROS production indicating that the observed differences arose primarily from events within the *bf* complex. These data demonstrate superoxide radical production in the cytochrome *bf* complex. They further support the hypothesis that blockage of the low potential chain promotes superoxide production by prolonging the lifetime of the reactive semiquinone generated at the quinol-oxidase (Q_o) site.

ACKNOWLEDGMENTS

Supported by NSF MCB 0091415, UW Oshkosh Faculty Development, and Centre National de la Recherche Scientifique, France. We thank Bill Antholine for generous assistance with EPR spectroscopy and Jiuseng Yan and Bill Cramer for kindly providing PetC–Δ2G.

REFERENCES

- Baymann, F., Rappaport, F., Joliot, P. & Kallas, T. (2001) *Biochemistry* 40: 10570–10577.
- Choquet, Y., Zito, F., Wostrikoff, K. & Wollman, F. A. (2003) *Plant Cell* 15: 1445–1454.
- Couture, M. M., Colbert, C. L., Babini, E., Rosell, F. I., Mauk, A. G., Bolin, J. T. & Eltis, L. D. (2001) *Biochemistry* 40: 84–92.
- de Vitry, C., Finazzi, G., Baymann, F. & Kallas, T. (1999) *Plant Cell* 11: 2031–2044.

- de Vitry, C., Ouyang, Y., Finazzi, G., Wollman, F. A. & Kallas, T. (2004) *J. Biol. Chem.* (in press).
- Gray, K. A., Dutton, P. L. & Daldal, F. (1994) *Biochemistry* 33: 723–733.
- Holton, B., Wu, X., Tsapin, A. I., Kramer, D. M., Malkin, R. & Kallas, T. (1996) *Biochemistry* 35: 15485–15493.
- Kurisu, G., Zhang, H., Smith, J. L. & Cramer, W. A. (2003) *Science* 302: 1009–1014.
- Lee, T. X., Metzger, S. U., Cho, Y. S., Whitmarsh, J. & Kallas, T. (2001) *Biochim. Biophys. Acta* 1504: 235–247.
- Merbitz-Zahradnik, T., Zwicker, K., Nett, J. N., Link, T. A. & Trumpower, B. L. (2003) *Biochemistry* 42: 13637–13645.
- Soriano, G. M., Guo, L. W., de Vitry, C., Kallas, T. & Cramer, W. A. (2002) *J. Biol. Chem.* 277: 1865–1871.
- Stroebel, D., Choquet, Y., Popot, J. L. & Picot, D. (2003) *Nature* 426: 413–418.
- Sutherland, M. W. & Learmonth, B. A. (1997) *Free Radic. Res.* 27: 283–289.
- Yan, J. & Cramer, W. A. (2003) *J. Biol. Chem.* 278: 20925–20933.

CYTOCHROME *b*₆ IN REDOX REGULATION OF PHOTOSYNTHESIS AND CHLOROPLAST GENE EXPRESSION

Jörgen Ström¹, Sujith Puthiyaveetil¹, Jens Forsberg², John F. Allen¹. ¹Plant Biochemistry, Lund University Box 124, Lund 221 00, Sweden. ²Medical Biochemistry and Microbiology Uppsala Biomedical Center, Uppsala University Box 582, Uppsala 751 23, Sweden

Keywords: Cytochrome *b*₆*f*, redox regulation, protein phosphorylation, transcriptional control, state transitions

INTRODUCTION

Cytochrome *b*₆ is a component of the cytochrome *b*₆*f* complex of the chloroplast thylakoid membrane. The recently described structures of cytochrome *b*₆*f* (Kurisu et al 2003, Strobel et al 2003) have provided new insight to the function of the complex (Allen 2004). We suggest that cytochrome *b*₆, apart from being important to both linear and cyclic electron transport, holds another key function, namely signaling in redox regulation of chloroplast function. This suggestion is supported by the observation that cytochrome *b*₆ is phosphorylated (Gal et al 1992) and by work of other laboratories showing that the occupancy of the Q_o quinol-binding site of the cytochrome *b*₆*f* complex is a factor governing the activation of a thylakoid protein kinase. Sequence comparisons reveal homology of an N-terminal segment of cytochrome *b*₆ with highly conserved phosphorylatable histidine residue, which sensor histidine kinases and receptor phytochromes could act upon (Hwang et al 2002, Grebe & Stock 1999). Recent phylogenetic analysis of Xiong and Bauer (2002) suggests that cytochrome *b*₆ is homologous with photosynthetic reaction centres, where the non-haem iron of type II centres (e.g., photosystem II) is coordinated by amino-acid side chains originally ligating haem. Electron transfer events in photosynthesis have been of over-riding importance in cell evolution, and in the function of the chloroplast genetic system.

MATERIALS AND METHODS

Chloroplast isolation and blue-native polyacrylamide gel electrophoresis. Leaves were collected from 12 day-old peas (*Pisum sativum*). The leaves were homogenised in grinding buffer (0.33 M sorbitol, 50 mM HEPES pH 7.6, 5 mM MgCl₂, 10 mM NaCl, 2 mM EDTA) and the homogenate was filtered through 2 layers, and then 8 layers, of cheese cloth. The filtrate was then layered onto a Percoll cushion and centrifuged at 600g for 7 min. Intact chloroplasts collected as a green pellet in the bottom of the tube were then resuspended in grinding buffer. The chloroplast suspension (0.33 mg Chl/ml) was incubated for 5 min in light with 400 μ Ci ³²P (Amersham) present. To the ³²P incubated chloroplast suspension was ferricyanide or dithiothreitol DTT (final concentration 2 mM), added, giving the oxidized and reduced chloroplast samples, respectively. These samples were transferred to darkness for 15 min. The remainder of the preparation was carried out under faint green light. To the chemically redox-treated chloroplasts NaF was added to give a final conc of 10 mM. This concentration of NaF was maintained for the remainder of the preparation. Chloroplasts were collected twice with Eppendorf microcentrifuge at 4000 rpm for 5 min, being resuspended and washed once with grinding buffer. The proteins in the final chloroplast preparation were separated using blue-native polyacrylamide gel electrophoresis essentially according to Kügler et al (1997).

Mass spectrometry analysis. Tryptic digests of excised protein spots were analysed with MALDI-TOF MS (Ultraflex, Bruker Daltonics). Protein identification was achieved by peptide mass fingerprinting.

Sequence comparison. The following databases and sequencing programs were used to establish the conserved N-terminal sequence of cyt *b₆*. Sequences aligned were downloaded from Swissprot Database. Multiple Alignment program used was ClustalW (<http://www.ebi.ac.uk/clustalw/>).

RESULTS

The Thylakoid Protein Complexes Show Different Levels of Phosphorylation. Figure 1A and 1B show blue-native polyacrylamide gels of thylakoid membranes reduced (A) and oxidized (B).




Figure 1: Coomassie stain and Phosphorimage of ³²P-labelled protein complexes from a blue native gel of redox treated chloroplasts. (A) and (B) are Coomassie-stained gels, reduced and oxidized. (C) and (D) are phosphorimages, reduced and oxidized. The numbered complex is Rubisco (1), ATPase/PSII (2), Cytochrome *b₆f* (3), PSII (4), LHCII (5).

Figure 1C and 1D shows the same gels, visualizing the ³²P labeled complexes by autoradiography.

The blue-native polyacrylamide gels are gradient gels (6–12% acrylamide) run without markers. The different protein complexes were identified using MALDI-TOF MS.

Phosphorylation of thylakoid complexes under oxidizing conditions. Figure 1D shows phosphorylation under oxidizing conditions. The major band in the autoradiograph of blue-native polyacrylamide gel representing the Rubisco (1) complex is strongly phosphorylated. The second most conspicuous band on the gel representing a mixture of PSII and ATPsynthase (2) is also strongly phosphorylated. The region of the gel corresponding to the *b₆f*-complex (3) has very low levels of phosphorylation, and the same is true for the second band, which represents PSII (4) proteins (CP47, D1, D2, cytochrome *b₅₅₉*). Finally the band containing LHCII protein (band 5, (containing a mixture of proteins, mostly Lhcb1, Lhcb2 and CP29)) show high levels of phosphorylations.

Phosphorylation of thylakoid complexes under reducing conditions. Figure 1C shows phosphorylation under reducing conditions, the major band in the blue-native polyacrylamide gel representing the Rubisco complex is phosphorylated to lesser extent than in Fig. 1D. The second band on the gel, representing a mixture of PSII and ATPsynthase, shows almost no phosphorylation. In contrast, the region of the gel corresponding to the *b₆f*-complex is now strongly phosphorylated. The second band, of PSII proteins (CP47, D1, D2, cytochrome *b₅₅₉*) shows the same low level of phosphorylation as under oxidizing conditions. Finally, the band containing LHCII proteins shows decreased phosphorylation under reducing conditions, but were still strongly phosphorylated.

Sequence Comparison of the N-terminal Region of *b₆*. We have aligned predicted gene products from *petB* (Fig. 2). The sequences aligned are N-terminal stretch of cyt *b₆*, length approximately 120 amino acids. A conserved histidine residue lies at the N-terminal part of the alignment (highlighted with a arrow). In the alignment, two of the four histidines, which ligate haem groups) are also seen. The other two haem-ligating histidines lie in the C-terminal part (not included in the alignment).

DISCUSSION

Phosphorylation of the Cytochrome *b₆f*-Complex. Cytochrome *b₆f* is involved in redox regulation in the chloroplast, being sensitive to changes in electron flow through the thylakoid membranes. We find that the cytochrome *b₆f* complex is strongly phosphorylated under reducing conditions. It is known that subunit V of the cytochrome *b₆f*-complex undergoes reversible phosphorylation (Hamel et al 2000) but subunit V seems to be present only in stroma lamella and not grana (Romanowska & Albertsson 1994). The redox state of the *b₆f*-complex acts as activator of the redox dependent LHCII kinase under reducing conditions (Aro & Ohad 2003). We think that cytochrome *b₆f* is closely involved in both activation of kinases and in regulation of RNA-synthesis. We suggest that the *b₆* protein in the complex is phosphorylated (Gal et al 1992) under reducing conditions. This phosphorylation might be the signal initiating changes in rates of RNA synthesis. How this is done is still unclear, but the conserved histidine at the N-terminus of *b₆* is a candidate for starting this signal.




Figure 2: Sequence alignment of cytochromes *b*₆ from 20 different species (*Chlamydomonas*, *Arabidopsis*, *Pisum*, *Physcomitrella*, *Nostoc*, *Synechocystis*, *Pinus* and more) showing the conserved histidine near the N-terminus.

ACKNOWLEDGMENTS

This work was supported by a Swedish Research Council grant to JFA and by Wallenberg Consortium North (WCN) through the Proteomics Resource Centre at Uppsala University.

REFERENCES

Allen, J. F. (2003a) *Philos. Trans. R. Soc.Lond. B* 358: 19–38.
 Allen, J. F. (2003b) *Comp. Funct. Genomics* vol. 4: 31–36.
 Allen, J. F. (2004) *Trends in Plant Sci.* vol. 3: 130–137.
 Aro, E. & Ohad, I. (2003) *Antioxidants & Redox Signaling* vol. 5 (nr 1): 55–67.
 Gal, A., Herrman, R. G., Lottspeich, F. & Ohad, I. (1992) *FEBS Lett.* 298: 33–35.
 Grebe, T. W. & Stock, J. B. (1999) *Advances in Micro. Physiol.* 41: 139–214.
 Hamel, P., Olive, J., Pierre, Y., Wollman, F. A. & de Vitry, C. A. (2000) *J. Biol. Chem.* 275: 17072–17079.
 Hwang, I., Chen, H. C. & Sheen, J. (2002) *Plant Physiol.* 129: 500–515.
 Kurisu, G., Zang, H., Smith, J. L. & Cramer, A. W. (2003) *Science* 302: 1009–1014.
 Kügler, M., Jansch, L., Kruff, V., Schmitz, U. K. & Braun, H. (1997) *Photosynthesis Res.* 53: 35–44.

Romanowska, E. & Albertsson, P. (1994) *Plant Cell Physiol.* 35: 557–568.
 Strobel, D., Choquet, Y., Popot, J. & Picot, D. (2003) *Nature* 426: 413–418.
 Xiong, J. & Bauer, C. E. (2002) *J. Mol. Biol.* 322: 1025–1037.

MOLECULAR BASIS OF THE MAJOR DIFFERENCE BETWEEN THE X-RAY STRUCTURE OF CYTOCHROME *b*₆*f* COMPLEX FROM *M. LAMINOSUS* AND *C. REINHARDTII*: THE BINDING SITE OF TRIDECYL-STIGMATELLIN

Jiusheng Yan, William A. Cramer. Department of Biological Sciences, Purdue University, West Lafayette, Indiana 47907-2054

Keywords: Cytochrome *b*₆*f* complex, Rieske iron-sulfur protein, stigmatellin, *Synechococcus* sp. PCC 7002

INTRODUCTION

The structures of the dimeric *b*₆*f* complex from the thermophilic cyanobacterium, *Mastigocladus laminosus* (Kurisu et al 2003), and the green alga, *Chlamydomonas reinhardtii* (Stroebel et al 2003), in the